



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (51) International Patent Classification ⁶ : C07D 235/08, A61K 31/415 | A1 | (11) International Publication Number: WO 97/47603 (43) International Publication Date: 18 December 1997 (18.12.97) |
| (21) International Application Number: PCT/SE97/00991 (22) International Filing Date: 5 June 1997 (05.06.97) (30) Priority Data: 9602286-8 10 June 1996 (10.06.96) SE (71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): AMIN, Kosrat [SE/SE]; Knarrhögsgatan 10 B, S-431 60 Mölndal (SE). DAHLSTRÖM, Mikael [SE/SE]; Marielundsgatan 5, S-431 67 Mölndal (SE). NORDBERG, Peter [SE/SE]; Björkebergsvägen 9, S-433 60 Sävedalen (SE). STARKE, Ingemar [SE/SE]; Frigångsgatan 8 C, S-413 01 Göteborg (SE). (74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE). | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: NEW COMPOUNDS (57) Abstract <p>The present invention relates to benzimidazole deriva- tives of formula (I), in which the phenyl moiety is substi- tuted with lower alkyl in 2- and 6-position, which inhibit exogenously or endogenously stimulated gastric acid secre- tion and thus can be used in the prevention and treatment of gastrointestinal inflammatory diseases.</p> <div style="text-align: center;"> </div> <div style="text-align: right;"> (I) </div> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|------------------------------------------|----|----------------------------------------------|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LJ | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

NEW COMPOUNDS

TECHNICAL FIELD


5 The present invention relates to novel compounds, and therapeutically acceptable salts thereof, which inhibit exogenously or endogenously stimulated gastric acid secretion and thus can be used in the prevention and treatment of gastrointestinal inflammatory diseases. In further aspects, the invention relates to compounds of the invention for use in therapy; to processes for preparation of such new
 10 compounds; to pharmaceutical compositions containing at least one compound of the invention, or a therapeutically acceptable salt thereof, as active ingredient; and to the use of the active compounds in the manufacture of medicaments for the medical use indicated above.

15


BACKGROUND ART

Benzimidazole derivatives of the following formula, active as anti-ulcer agents, are disclosed in EP-B-0 266 326 and in US 5,106,862:

20



wherein R^1 is i.a. an aryl group of the formula



in which each of R^7 , R^8 and R^9 independently represents i.a. H or C_1 - C_6 alkyl, or halogen;

5 R^2 is i.a. H;

R^3 is i.a. H or C_1 - C_6 alkyl;

n is an integer 0-6;

R^4 , R^5 and R^6 are H or C_1 - C_6 alkyl;

A is alkylene up to 6 carbon atoms optionally interrupted by a hetero atom such as

10 O or N.


For a review of the pharmacology of the gastric acid pump (the H^+ , K^+ -ATPase), see Sachs et al. (1995) Annu. Rev. Pharmacol. Toxicol. 35: 277-305.

15

DISCLOSURE OF THE INVENTION

It has surprisingly been found that compounds of the Formula I, which are substituted benzimidazole derivatives in which the phenyl moiety is substituted
20 with lower alkyl in 2- and 6-position, are particularly effective as inhibitors of the gastrointestinal H^+ , K^+ -ATPase and thereby as inhibitors of gastric acid secretion.

In one aspect, the invention thus relates to compounds of the general Formula I:



I

or a pharmaceutically acceptable salt thereof, wherein

5 R¹ is lower alkyl;

R² is lower alkyl;

R³, which is in position 3, 4, or 5 of the phenyl ring, is

- 10 (a) H,
(b) halogen, or
(c) lower alkyl;

R⁴ is

- 15 (a) H, or
(b) lower alkyl;

X, which is connected to the heterocycle in the position 4 or 7, is

- 20 (a) NH, or
(b) O.

As used herein, the term "lower alkyl" denotes a straight or branched alkyl group having from 1 to 6 carbon atoms. Examples of said lower alkyl include methyl,

ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl and straight- and branched-chain pentyl and hexyl.

The term "halogen" includes fluoro, chloro, bromo and iodo.

5

Both pure enantiomers, racemic mixtures and unequal mixtures of two enantiomers are within the scope of the invention. It should be understood that all the diastereomeric forms possible (pure enantiomers, racemic mixtures and unequal mixtures of two enantiomers) are within the scope of the invention. Also
10 included in the invention are derivatives of the compounds of the Formula I which have the biological function of the compounds of the Formula I.

Depending on the process conditions the end products of the Formula I are obtained either in neutral or salt form. Both the free base and the salts of these end
15 products are within the scope of the invention.

Acid addition salts of the new compounds may in a manner known *per se* be transformed into the free base using basic agents such as alkali or by ion exchange. The free base obtained may also form salts with organic or inorganic
20 acids.

In the preparation of acid addition salts, preferably such acids are used which form suitably therapeutically acceptable salts. Examples of such acids are hydrohalogen acids such as hydrochloric acid, sulfuric acid, phosphoric acid,
25 nitric acid, aliphatic, alicyclic, aromatic or heterocyclic carboxyl or sulfonic acids, such as formic acid, acetic acid, propionic acid, succinic acid, glycolic acid, lactic acid, malic acid, tartaric acid, citric acid, ascorbic acid, maleic acid, hydroxymaleic acid, pyruvic acid, p-hydroxybenzoic acid, embonic acid, methanesulfonic acid, ethanesulfonic acid, hydroxyethanesulfonic acid, halogenbensenesulfonic acid,
30 toluenesulfonic acid or naphthalenesulfonic acid.

Preferred compounds according to the invention are those of the formula I wherein R¹ is CH₃ or CH₂CH₃; R² is CH₃ or CH₂CH₃; R³ is H, 4-F or 4-Cl; R⁴ is H or CH₃ and X is NH or O connected to the heterocycle in the 4-position.

5

Particularly preferred compounds according to the invention are:

4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole;

4-(2,6-dimethylbenzyloxy)-2-methylbenzimidazole;

4-(2,6-dimethyl-4-fluorobenzylamino)-2-methylbenzimidazole;

10 4-(2,6-dimethyl-4-fluorobenzyloxy)-2-methylbenzimidazole;

4-(2,6-dimethylbenzylamino)-1,2-dimethylbenzimidazole;

4-(2-ethyl-6-methylbenzylamino)-2-methylbenzimidazole;

4-(2,6-diethylbenzylamino)-2-methylbenzimidazole;

4-(2,6-dimethyl-4-fluorobenzylamino)-1,2-dimethylbenzimidazole;

15 4-(2,6-dimethyl-4-fluorobenzyloxy)-1,2-dimethylbenzimidazole.

Preparation

The present invention also provides the following processes A to C for the manufacture of compounds with the general Formula I:


20

Process A

Process A for manufacture of compounds with the general Formula I comprises the following steps:

25


a) Compounds of the general Formula II



II

wherein X^1 is NH_2 or OH connected to the heterocycle in the position 4 or 7 and R_4 is as defined for Formula I, can be reacted with compounds of the general

5 Formula III



III


wherein R^1 , R^2 and R^3 are as defined for Formula I and Y is a leaving group, such
 10 as a halide, tosyloxy or mesyloxy, to the compounds of the Formula I.

It is convenient to conduct this reaction in an inert solvent, e.g. acetone, acetonitrile, dimethoxyethane, methanol, ethanol or dimethylformamide with or without a base. The base is e.g. an alkali metal hydroxide, such as sodium
 15 hydroxide and potassium hydroxide; a sodium alcoholate, such as sodium methoxide and sodium ethoxide; an alkali metal hydride such as sodium hydride and potassium hydride; an alkali metal carbonate, such as potassium carbonate and sodium carbonate; or an organic amine, such as triethylamin.


20 Process B

Process B for manufacture of compounds with the general Formula I comprises the following step:

A compound of the Formula IV




- 5 wherein R^4 is as defined for Formula I and the NH_2 group is connected to the heterocycle in the position 4 or 7, can be reacted with compounds of the general Formula V



10

- wherein R^1 , R^2 and R^3 are as defined for Formula I, in the presence of a Lewis acid e.g. zinc chloride to the compounds of the Formula VI



15

- wherein R^4 is as defined for Formula I and the imine nitrogen is connected to the heterocycle in the position 4 or 7, whereby the compounds of the general Formula

VI are reduced e.g. by using sodium borhydride or sodiumcyano borhydride to compounds of the general Formula I. The reactions can be carried out under standard conditions in an inert solvent e.g. methanol or ethanol.

5 *Process C*

Compounds of the general Formula I wherein R⁴ is "lower alkyl" can be made by alkylation of compounds of the general Formula I wherein R⁴ is H in an inert solvent, e.g. acetonitrile or acetone, with or without base with compounds of the
10 general Formula VII



VII

wherein R⁴ is as defined for Formula I and X² is a leaving group e.g. a halide,
15 mesylate or tosylate. The base is e.g. an alkali metal hydroxide, such as sodium hydroxide and potassium hydroxide; a sodium alcoholate, such as sodium methoxide and sodium ethoxide; an alkali metal hydride such as sodium hydride and potassium hydride; an alkali metal carbonate, such as potassium carbonate and sodium carbonate; or an organic amine, such as triethylamine.

20

Medical use

In a further aspect, the invention relates to compounds of the formula I for use in therapy, in particular for use against gastrointestinal inflammatory diseases. The
25 invention also provides the use of a compound of the formula I in the manufacture of a medicament for the inhibition of gastric acid secretion, or for the treatment of gastrointestinal inflammatory diseases.

The compounds according to the invention may thus be used for prevention and
30 treatment of gastrointestinal inflammatory diseases, and gastric acid-related

diseases in mammals including man, such as gastritis, gastric ulcer, duodenal ulcer, reflux esophagitis and Zollinger-Ellison syndrom. Furthermore, the compounds may be used for treatment of other gastrointestinal disorders where gastric antisecretory effect is desirable, e.g. in patients with gastrinomas, and in
5 patients with acute upper gastrointestinal bleeding. They may also be used in patients in intensive care situations, and pre-and postoperatively to prevent acid aspiration and stress ulceration.

Pharmaceutical formulations

10

In yet a further aspect, the invention relates to pharmaceutical compositions containing at least one compound of the invention, or a therapeutically acceptable salt thereof, as active ingredient.

15 The compounds of the invention can also be used in formulations together with other active ingredients, e.g. antibiotics such as amoxicillin.

For clinical use, the compounds of the invention are formulated into pharmaceutical formulations for oral, rectal, parenteral or other mode of
20 administration. The pharmaceutical formulation contains a compound of the invention in combination with one or more pharmaceutically acceptable ingredients. The carrier may be in the form of a solid, semi-solid or liquid diluent, or a capsule. These pharmaceutical preparations are a further object of the invention. Usually the amount of active compounds is between 0.1–95% by weight
25 of the preparation, preferably between 0.1–20% by weight in preparations for parenteral use and preferably between 0.1 and 50% by weight in preparations for oral administration.

In the preparation of pharmaceutical formulations containing a compound of the
30 present invention in the form of dosage units for oral administration the

compound selected may be mixed with solid, powdered ingredients, such as lactose, saccharose, sorbitol, mannitol, starch, amylopectin, cellulose derivatives, gelatin, or another suitable ingredient, as well as with disintegrating agents and lubricating agents such as magnesium stearate, calcium stearate, sodium stearyl fumarate and polyethylene glycol waxes. The mixture is then processed into granules or pressed into tablets.

Soft gelatine capsules may be prepared with capsules containing a mixture of the active compound or compounds of the invention, vegetable oil, fat, or other suitable vehicle for soft gelatine capsules. Hard gelatine capsules may contain granules of the active compound. Hard gelatine capsules may also contain the active compound in combination with solid powdered ingredients such as lactose, saccharose, sorbitol, mannitol, potato starch, corn starch, amylopectin, cellulose derivatives or gelatine.

Dosage units for rectal administration may be prepared (i) in the form of suppositories which contain the active substance mixed with a neutral fat base; (ii) in the form of a gelatine rectal capsule which contains the active substance in a mixture with a vegetable oil, paraffin oil or other suitable vehicle for gelatine rectal capsules; (iii) in the form of a ready-made micro enema; or (iv) in the form of a dry micro enema formulation to be reconstituted in a suitable solvent just prior to administration.

Liquid preparations for oral administration may be prepared in the form of syrups or suspensions, e.g. solutions or suspensions containing from 0.1% to 20% by weight of the active ingredient and the remainder consisting of sugar or sugar alcohols and a mixture of ethanol, water, glycerol, propylene glycol and polyethylene glycol. If desired, such liquid preparations may contain colouring agents, flavouring agents, saccharine and carboxymethyl cellulose or other thickening agent. Liquid preparations for oral administration may also be

prepared in the form of a dry powder to be reconstituted with a suitable solvent prior to use.

Solutions for parenteral administration may be prepared as a solution of a compound of the invention in a pharmaceutically acceptable solvent, preferably in a concentration from 0.1% to 10% by weight. These solutions may also contain stabilizing ingredients and/or buffering ingredients and are dispensed into unit doses in the form of ampoules or vials. Solutions for parenteral administration may also be prepared as a dry preparation to be reconstituted with a suitable solvent extemporaneously before use.

The typical daily dose of the active substance varies within a wide range and will depend on various factors such as for example the individual requirement of each patient, the route of administration and the disease. In general, oral and parenteral dosages will be in the range of 5 to 1000 mg per day of active substance.


The compounds according to the invention can also be used in formulations together with other active ingredients, e.g. for the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa. Such other active ingredients may be antimicrobial agents, in particular:

- b-lactam antibiotics such as amoxicillin, ampicillin, cephalothin, cefaclor or cefixime;
- macrolides such as erythromycin, or clarithromycin;
- tetracyclines such as tetracycline or doxycycline;
- aminoglycosides such as gentamycin, kanamycin or amikacin;
- quinolones such as norfloxacin, ciprofloxacin or enoxacin;
- others such as metronidazole, nitrofurantoin or chloramphenicol; or
- preparations containing bismuth salts such as bismuth subcitrate, bismuth subsalicylate, bismuth subcarbonate, bismuth subnitrate or bismuth subgallate.

EXAMPLES

1. PREPARATION OF COMPOUNDS OF THE INVENTION


5

*Example 1.1**Synthesis of 4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole*

10

4-amino-2-methylbenzimidazole (6.0 g, 0.041 mol) was dissolved in acetonitrile (120 ml). To the solution were added 2,6-dimethylbenzylchloride (6.3 g, 0.041 mol), sodium carbonate (16 g, 0.15 mol) and a cat. amount of sodium iodide and the reaction mixture was refluxed for 3 h. The sodium carbonate was removed by
15 filtration and was washed with methylene chloride. Vacuum evaporation of the solvent gave an oily residue which was subjected to flash chromatography on silica gel, methylene chloride : methanol (10:1) . The residue was crystallized by treating with ethyl acetate to obtain 3.3 g (30 %) of the title compound.

20 ¹H-NMR, 300 MHz, CDCl₃): 2.35 (s, 6H), 2.50 (s, 3H), 4.35 (s, 2H), 4.45 (bs, 1H), 6.50 (d, 1H), 6.75 (d, 1H), 6.95-7.15 (m, 4H)


*Example 1.2**Synthesis of 4-(2,6-dimethylbenzyloxy)-2-methylbenzimidazole*

5

4-hydroxy-2-methylbenzimidazole (0.59 g, 4 mmol) was dissolved in acetonitrile (15 ml). To the solution were added 2,6-dimethylbenzylchloride (0.52 g, 4 mmol) and sodium hydroxide (0.16 g, 4 mmol) (dissolved in 1 ml water) and the reaction mixture was refluxed for 2 h. The solvent was evaporated under reduced pressure and the residue was dissolved in methylene chloride and was washed with 2 M NaOH. The organic layer was separated, dried over sodium sulfate and evaporated under reduced pressure. The oily residue was purified by column chromatography on silica gel. The product was eluted with methylene chloride containing 5 % methanol. Crystallization from acetonitrile gave 0.18 g (18 %) of the title compound.

15

(¹H-NMR, 300 MHz, DMSO-d₆): 2.3 (s, 6H), 2.40 (s, 3H), 5.25 (s, 2H), 6.9 (d, 1H), 7.05-7.15 (m, 4H), 7.2 (t, 1H)


*Example 1.3**Synthesis of 4-(2,6-dimethyl-4-fluorobenzylamino)-2-methylbenzimidazole*

5

4-amino-2-methylbenzimidazole (0.6 g, 4.1 mmol) was dissolved in acetonitrile (10 ml). To the solution were added 2,6-dimethyl-4-fluoro-benzylbenzylbromide (0.89 g, 4.1 mol) and potassium carbonate (0.68 g, 4.9 mmol) and the mixture was warmed at 70-80 °C for 3 h. The reaction mixture was cooled to room temperature and methylene chloride (25 ml) and water (25 ml) was added. The organic layer was separated, dried over sodium sulfate and was evaporated under reduced pressure. The residue was crystallized twice from ethyl acetate and was purified by column chromatography on silica gel, using methylene chloride containing 10% methanol as eluent. There was obtained 0.26 g (22 %) of the title compound.

15


(¹H-NMR, 300 MHz, CDCl₃): 2.3 (s, 6H), 2.45 (s, 3H), 4.25 (s, 2H), 4.35 (bs, 1H), 6.45 (d, 1H), 6.65 (d, 2H), 6.7 (d, 1H), 7.1 (t, 1H)

*Example 1.4**Synthesis of 4-(2,6-dimethyl-4-fluorobenzoyloxy)-2-methylbenzimidazole*

5

4-hydroxy -2-methylbenzimidazole (0.57 g, 3.8 mmol) was dissolved in acetonitrile (4 ml). To the solution were added sodium hydroxide (0.17 g, 43 mmol) (dissolved in 1 ml water) and 2,6-dimethyl-4-fluorobenzylbromide (0.82 g, 3.8 mmol) dissolved in acetonitrile (6 ml) and the reaction mixture was refluxed
10 for 80 min. and was stirred for 20 h. at ambient temperature. To the reaction mixture was added methylene chloride (25 ml) and water (25 ml). The organic layer was separated, washed three times with 2 M NaOH, dried over sodium sulfate and was evaporated under reduced pressure. The residue was crystallized from ethyl acetate and was then purified twice by column chromatography on
15 silica gel a) methylene chloride : methanol (10:1) b) ethyl acetate : methanol : acetic acid : water (96:6:6:4). There was obtained 0.066 g (6 %) of the title compound.


(¹H-NMR, 300 MHz, DMSO-d₆): 2.3 (s, 6H), 2.5 (s, 3H), 5.05 (s, 2H), 6.7 (d, 2H),
20 6.8 (d, 1H), 7.15 (t, 1H), 7.25 (d, 1H)

*Example 1.5**Synthesis of 4-(2,6-dimethylbenzylamino)-1,2-dimethylbenzimidazole*

5

4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole (0.2 g 0.75 mmol) was dissolved in acetonitrile (15 ml). To the solution were added methyl iodide (0.05 ml 0.82 mmol) and potassium carbonate (0.2 g 1.4 mmol) and the reaction mixture was stirred at ambient temperature for 20 h. The solids were removed by filtration and was washed with methylene chloride. Vacuum evaporation of the solvent
10 gave an oily residue which was subjected to flash chromatography on silica gel, methylene chloride : methanol (10:1) . Crystallization from acetonitrile gave 0.018 g (9%) of the title compound.

15 (¹H-NMR, 500 MHz, CDCl₃): 2.4 (s, 6H), 2.55 (s, 3H), 3.65 (s, 3H), 4.4 (d, 2H), 4.55 (bs, 1H), 6.55 (d, 1H), 6.7 (d, 1H) 7.05 (d, 2H), 7.1 (t, 1H), 7.2 (t, 1H).


*Example 1.6**Synthesis of 4-(2-Ethyl-6-methylbenzylamino)-2-methylbenzimidazole*

5

4-amino-2-methylbenzimidazole (0.4 g, 2.7 mmol) and 2-ethyl-6-methylbenzylchloride (0.46 g, 2.7 mmol) were dissolved in 10 ml dimethoxyethane. Sodium carbonate (0.5 g, 4.7 mmol) and (0.2 g, 4.7 mmol) of potassium iodide were added to the solution. The reaction mixture was refluxed
10 for 2 h. Inorganic salts were removed by filtration and washed with dimethoxyethane. The filtrate was evaporated to dryness. The residue was purified by column chromatography on silica gel. The product was eluted with a 50:50 mixture of methylene chloride and ethyl acetate. There was obtained 0.21 g of the title compound.


15

$^1\text{H-NMR}$, 400 MHz, CDCl_3 : 1.1 (t, 3H), 2.35 (s, 3H), 2.45 (s, 3H), 2.70 (q, 2H), 4.35 (s, 2H), 4.45 (bs, 1H), 6.55 (d, 1H), 6.75 (d, 1H), 6.95-7.20 (m, 4H).

*Example 1.7**Synthesis of 4-(2,6-diethylbenzylamino)-2-methylbenzimidazole*

5 4-Amino-2-methylbenzimidazole (0.79 g, 5.4 mmol) and 2,6-diethylbenzaldehyde (1.1 g, 6.6 mmol) were dissolved in methanol (30 ml). ZnCl_2 (0.9 g, 6.6 mmol) and subsequently NaBH_3CN (0.42 g, 6.6 mmol) in small portions were added and the mixture was refluxed under argon for 3 hours and was stirred at room
10 temperature for 16 h. The mixture was poured over an aqueous 1M NaOH solution (50 ml). The resultant yellow suspension was extracted with DCM and the organic solution was washed with brine, dried over Na_2SO_4 and then evaporated under reduced pressure. The oily residue (1.8 g) was purified by crystallization from a mixture of ethyl acetate and acetonitrile to give 0.65g (41%)
15 of the title compound.

($^1\text{H-NMR}$, 300 MHz, CDCl_3): 1.2 (t, 6H), 2.5 (s, 3H), 2.27 (q, 4H), 4.35 (d, 2H), 4.55 (bs, 1H), 6.05 (d, 1H), 6.75 (d, 1H), 7.0-7.25 (m, 4H), 9.0 (bs, 1H)


*Example 1.8**Synthesis of 4-(2,6-dimethyl-4-fluorobenzylamino)-1,2-dimethylbenzimidazole.*

5

4-(2,6-Dimethyl-4-fluorobenzylamino)-2-methylbenzimidazole (0.1 g, 0.35 mmol) was dissolved in 1,2-dimethoxyethane (3 ml). Solid sodium hydroxide (25 mg, 0.63 mmol) and tetrabutylammonium bromide (5 mg, 0.016 mmol) were added. The mixture was stirred 15 minutes at ambient temperature. Methyl iodide (60 mg, 0.42 mmol) was added and the reaction mixture was stirred at ambient temperature for 2.5 h. The solvent was evaporated under reduced pressure. The residue was purified by column chromatography on silica gel, using methylene chloride : ethyl acetate (50 : 50) as eluent. 80 mg (76%) of the title compound was obtained.

15

¹H-NMR, 300 MHz, CDCl₃): 2.38 (s, 6H), 2.53 (s, 3H), 3.67 (s, 3H), 4.32 (d, 2H), 4.44 (bs, 1H), 6.51 (d, 1H), 6.68 (d, 1H), 6.75 (d, 2H), 7.18 (t, 1H).

*Example 1.9**Synthesis of 4-(2,6-dimethyl-4-fluorobenzyloxy)-1,2-dimethylbenzimidazole.*

5

4-(2,6-dimethyl-4-fluorobenzyloxy)-2-methylbenzimidazole (180 mg, 0.63 mmol) was dissolved in 1,2-dimethoxyethane (6 ml). To the solution was added sodium hydroxide (26 mg, 0.65 mmol) and methyl iodide (0.47 ml, 0.76 mmol). The reaction mixture was stirred at ambient temperature for 3.5 h. After evaporation of the solvent, the residue was purified by chromatography on silica gel, methylene chloride : methanol 10:1. Chromatography of one of the obtained fractions on silica using ethyl acetate as eluent gave 40 mg (21%) of the title compound.

15 ¹H-NMR, 300 MHz, CDCl₃): 2.40 (s, 6H), 2.58 (s, 3H), 3.70 (s, 3H), 5.24 (s, 2H), 6.75 (d, 2H), 6.82 (d, 1H), 6.93 (d, 1H), 7.19 (t, 1H).

Example 1.10

Preparation of the hydrochloride salt of 4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole

20

4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole (0.5g 1.9 mmol) was dissolved in ethyl acetate (20 ml) and methanol (6 ml). To the solution was added

hydrochloric acid (12 M) (0.16 ml, 1.9 mol) and the mixture was stirred at ambient temperature for 5 min. The precipitated salt was filtered off and dried to give the desired product as a white crystalline solid (0.45 g, 79%).

5 ($^1\text{H-NMR}$, 500 MHz, MeOD): 2.4 (s, 6H), 2.8 (s, 3H), 4.45 (s, 2H), 6.9 (d, 1H), 7.0 (d, 1H), 7.1 (d, 2H), 7.15 (t, 1H), 7.4 (t, 1H).

Example 1.11

Preparation of the methanesulfonic acid salt of 4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole
10 methylbenzimidazole

4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole (0.5 g, 1.9 mmol) was dissolved in a ethyl acetate (20 ml) and methanol (7 ml). To the solution was added methanesulfonic acid (0.18 g, 1.9 mmol) and the mixture was stirred at
15 ambient temperature for 5 min. The precipitated salt was filtered off and dried to give the desired product as a white crystalline solid (0.58 g, 85%).

($^1\text{H-NMR}$, 500 MHz, MeOD): 2.4 (s, 6H), 2.65 (s, 3H), 2.75 (s, 3H), 4.45 (s, 2H), 6.9 (d, 1H), 7.0 (d, 1H), 7.1 (d, 2H), 7.15 (t, 1H), 7.4 (t, 1H).

20

2. PREPARATION OF INTERMEDIATES

Example 2.1

25 *Synthesis of 2,6-dimethyl-4-fluoro-benzylbromide*

A mixture of 3,5-dimethyl-fluorobenzene (5 g, 0.04 mol), paraformaldehyde (15 g), hydrobromic acid (70 ml) (30 % in acetic acid) and acetic acid (25 ml) was stirred at ambient temperature for 4.5 h. To the mixture were water and
30 petroleum ether added and the organic layer was separated dried over anhydrous

sodium sulfate and evaporated carefully under reduced pressure. The residue was purified by column chromatography on silica gel with petroleum ether as eluent to give the title product. (3.7 g, 43 %)

5 (¹H-NMR, 300 MHz, CDCl₃): 2.5 (s, 6H), 4.55 (s, 2H), 6.75 (d, 2H)

Example 2.2

Synthesis of 2-ethyl-6-methylbenzylchloride

10 2-Ethyl-6-methylbenzylalcohol (1.0 g, 6.67 mmol) was dissolved in 10 ml methylene chloride. Thionyl chloride (1.0 g, 8.5 mmol) was added. The mixture was stirred over night at ambient temperature. The reaction mixture was evaporated. The residue was dissolved in methylene chloride and filtered through 5 g of silica gel. The filtrate was evaporated. 1.0 g (89 %) of the title compound
15 (oil) was obtained.

(¹H-NMR, 300 MHz, CDCl₃): 1.29 (t, 3H), 2.46 (s, 3H), 2.76 (q, 2H), 4.71 (s, 2H), 7.0-7.2 (m, 3H).

20

3. BIOLOGICAL TESTS

3.1. *In vitro* experiments

25 *Acid secretion inhibition in isolated rabbit gastric glands*

Inhibiting effect on acid secretion *in vitro* in isolated rabbit gastric glands was measured as described by Berglindh et al. (1976) Acta Physiol. Scand. 97, 401-414.

The compounds according to Example 1.1 to 1.7 exhibited IC₅₀ values of less than 1.6 μ M.

Compound (A) for comparison (4-benzylamino-2-methylbenzimidazole) is a compound as claimed in claim 1 of EP-B-0 266 326 wherein:


A is CH₂NH;

R², R³, R⁴, R⁵ and R⁶ are H;

n is 1; and

R¹ is a group of the formula II' wherein R⁷, R⁸ and R⁹ are H.

Compound (A) for comparison exhibited an IC₅₀ value of 12.9 μ M.



(A)

Compound (B) for comparison (4-benzoyloxy-2-methyl benzimidazole) is a compound as claimed in claim 1 of EP-B-0 266 326 wherein:


A is CH₂O;

R², R³, R⁴, R⁵ and R⁶ are H;

n is 1; and

R¹ is a group of the formula II' wherein R⁷, R⁸ and R⁹ are H.

Compound (B) for comparison exhibited an IC₅₀ value of 7.8 μ M.



(B)

Consequently, the compounds according to the invention exhibited a significantly better inhibiting effect on acid secretion *in vitro* compared to the compounds (A) and (B), which carry an unsubstituted phenyl group. It is concluded that the improved effect is due to the phenyl group, in the compounds according to the invention, being substituted with lower alkyl in 2- and 6-position.

3.2. *In vivo* experiments

Inhibiting effect on acid secretion in female rats

Female rats of the Sprague-Dawley strain were used. They were equipped with cannulated fistulae in the stomach (lumen) and the upper part of the duodenum, for collection of gastric secretions and administration of test substances, respectively. A recovery period of 14 days after surgery was allowed before testing commenced.

Before secretory tests, the animals are deprived of food but not water for 20 h. The stomach is repeatedly washed through the gastric cannula with tap water (37°C), and 6 ml Ringer-Glucose given subcutaneously. Acid secretion is stimulated with infusion during 2.5-4 h (1.2 ml/h, subcutaneously) of pentagastrin and carbachol (20 and 110 nmol/kg·h, respectively), during which time gastric secretions are collected in 30-min fractions. Test substances or vehicle are given either at 60 min

after starting the stimulation (intravenous and intraduodenal dosing, 1 ml/kg), or 2 h before starting the stimulation (oral dosing, 5 ml/kg, gastric cannula closed). The time interval between dosing and stimulation may be increased in order to study the duration of action. Gastric juice samples are titrated to pH 7.0 with NaOH, 0.1 M, and acid output calculated as the product of titrant volume and concentration.

Further calculations are based on group mean responses from 4-6 rats. In the case of administration during stimulation; the acid output during the periods after administration of test substance or vehicle are expressed as fractional responses, setting the acid output in the 30-min period preceding administration to 1.0. Percentage inhibition is calculated from the fractional responses elicited by test compound and vehicle. In the case of administration before stimulation; percentage inhibition is calculated directly from acid output recorded after test compound and vehicle.

Bioavailability in rat

Adult rats of the Sprague-Dawley strain were used. One to three days prior to the experiments all rats were prepared by cannulation of the left carotid artery under anaesthesia. The rats used for intravenous experiments were also cannulated in the jugular vein (Popovic (1960) J. Appl. Physiol. 15, 727-728). The cannulas were exteriorized at the nape of the neck.

Blood samples (0.1 - 0.4 g) were drawn repeatedly from the carotid artery at intervals up to 5.5 hours after given dose. The samples were frozen until analysis of the test compound.

Bioavailability was assessed by calculating the quotient between the area under blood/plasma concentration (AUC) curve following (i) intraduodenal (i.d.) or oral

(p.o.) administration and (ii) intravenous (i.v.) administration from the rat or the dog, respectively.

The area under the blood concentration vs. time curve, AUC, was determined by the log/linear trapezoidal rule and extrapolated to infinity by dividing the last
5 determined blood concentration by the elimination rate constant in the terminal phase. The systemic bioavailability (F%) following intraduodenal or oral administration was calculated as

$$F(\%) = (AUC \text{ (p.o. or i.d.)} / AUC \text{ (i.v.)}) \times 100.$$

10

Inhibition of gastric acid secretion and bioavailability in conscious dog

Labrador retriever or Harrier dogs of either sex were used. They were equipped with a duodenal fistula for the administration of test compounds or vehicle and a
15 cannulated gastric fistula or a Heidenhaim-pouch for the collection of gastric secretion.

Before secretory tests the animals were fasted for about 18 h but water was freely allowed. Gastric acid secretion was stimulated for up to 6.5 h infusion of
20 histamine dihydrochloride (12 ml/h) at a dose producing about 80% of the individual maximal secretory response, and gastric juice collected in consecutive 30-min fractions. Test substance or vehicle was given orally, i.d. or i.v., 1 or 1.5 h after starting the histamine infusion, in a volume of 0.5 ml/kg body weight. In the case of oral administration, it should be pointed out that the test compound is
25 administered to the acid secreting main stomach of the Heidenham-pouch dog.


The acidity of the gastric juice samples were determined by titration to pH 7.0, and the acid output calculated. The acid output in the collection periods after administration of test substance or vehicle were expressed as fractional responses,
30 setting the acid output in the fraction preceding administration to 1.0. Percentage

inhibition was calculated from fractional responses elicited by test compound and vehicle.

Blood samples for the analysis of test compound concentration in plasma were
5 taken at intervals up to 4 h after dosing. Plasma was separated and frozen within
30 min after collection and later analyzed. The systemic bioavailability (F%) after
oral or i.d. administration was calculated as described above in the rat model.

CLAIMS

1. A compound of the Formula I



I

or a pharmaceutically acceptable salt thereof, wherein

R¹ is lower alkyl;

R² is lower alkyl;

R³, which is in position 3, 4, or 5 of the phenyl ring, is

(a) H,

(b) halogen, or

(c) lower alkyl;

R⁴ is

(a) H, or

(b) lower alkyl;

X, which is connected to the heterocycle in the position 4 or 7, is


(a) NH, or

(b) O.

2. A compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein X is connected to the heterocycle in position 4.
- 5 3. A compound according to claim 2, or a pharmaceutically acceptable salt thereof, wherein R^1 is CH_3 or CH_2CH_3 ; R^2 is CH_3 or CH_2CH_3 ; R^3 is H, 4-F or 4-Cl; and R^4 is H or CH_3 .
- 10 4. A compound according to claim 3 which is the compound 4-(2,6-dimethylbenzylamino)-2-methylbenzimidazole, or a pharmaceutically acceptable salt thereof.
- 15 5. A compound according to claim 3 which is the compound 4-(2,6-dimethylbenzyloxy)-2-methylbenzimidazole, or a pharmaceutically acceptable salt thereof.
- 20 6. A compound according to claim 3 which is the compound 4-(2,6-dimethyl-4-fluorobenzylamino)-2-methylbenzimidazole, or a pharmaceutically acceptable salt thereof.
- 25 7. A compound according to claim 3 which is the compound 4-(2,6-dimethyl-4-fluorobenzyloxy)-2-methylbenzimidazole, or a pharmaceutically acceptable salt thereof.
8. A compound according to claim 3 which is the compound 4-(2,6-dimethylbenzylamino)-1,2-dimethylbenzimidazole, or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 3 which is the compound 4-(2-ethyl-6-methylbenzylamino)-2-methylbenzimidazole, or a pharmaceutically acceptable salt thereof.
- 5 10. A compound according to claim 3 which is the compound 4-(2,6-diethylbenzylamino)-2-methylbenzimidazole, or a pharmaceutically acceptable salt thereof.
- 10 11. A compound according to claim 3 which is the compound 4-(2,6-dimethyl-4-fluorobenzylamino)-1,2-dimethylbenzimidazole, or a pharmaceutically acceptable salt thereof.
- 15 12. A compound according to claim 3 which is the compound 4-(2,6-dimethyl-4-fluorobenzyloxy)-1,2-dimethylbenzimidazole, or a pharmaceutically acceptable salt thereof.
13. A compound which is a hydrochloride salt of a compound according to any one of claims 1 to 12.
- 20 14. A compound which is a methanesulfonic acid salt of a compound according to any one of claims 1 to 12.
15. A compound according to any one of claims 1 to 14 for use in therapy.
- 25 16. A process for the preparation of a compound according to any one of claims 1 to 14 comprising reacting a compound of the general Formula II


31



II

wherein X^1 is NH_2 or OH connected to the heterocycle in the position 4 or 7 and R^4 is as defined for Formula I, with a compound of the general Formula

III




III

wherein R^1 , R^2 and R^3 are as defined for Formula I and Y is a leaving group, in an inert solvent with or without a base, to a compound of the Formula I.

17. A process for the preparation of a compound according to any one of claims 1 to 14 comprising


(a) reacting a compound of the Formula IV



IV

wherein R^4 is as defined for Formula I and the NH_2 group is connected to the heterocycle in the position 4 or 7, with a compound of the Formula V


32



V

wherein R^1 , R^2 and R^3 are as defined for Formula I, in the presence of a Lewis acid in an inert solvent to a compound of the Formula VI

5



VI

wherein R^4 is as defined for Formula I and the imine nitrogen is connected to the heterocycle in the position 4 or 7.

10

(b) reducing a compound of the Formula VI in an inert solvent under standard conditions to a compound of the general Formula I.

18. A process for the preparation of a compound according to any one of claims 1, 2, 3 or 8 wherein R^4 is "lower alkyl" comprising

15

alkylation of a compound of the Formula I wherein R^4 is H in an inert solvent with or without base with a compound of the Formula VII

20



VII

wherein R^4 is as defined for Formula I and X^2 is a leaving group, to a compound of the Formula I wherein R^4 is "lower alkyl".

- 5 19. A pharmaceutical formulation comprising a compound according to any one of claims 1 to 14 and in addition comprising a pharmaceutically acceptable carrier.
20. Use of a compound according to any one of claims 1 to 14 for the manufacture
10 of a medicament for the inhibition of gastric acid secretion.
21. Use of a compound according to any one of claims 1 to 14 for the manufacture of a medicament for the treatment of gastrointestinal inflammatory diseases.
- 15 22. Use of a compound according to any one of claims 1 to 14 for the manufacture of a medicament for the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa, wherein the said salt is adapted to be administered in combination with at least one antimicrobial agent.
- 20 23. A method for inhibiting gastric acid secretion which comprises administering to a mammal, including man, in need of such inhibition an effective amount of a compound according to any one of claims 1 to 14.
- 25 24. A method for the treatment of gastrointestinal inflammatory diseases which comprises administering to a mammal, including man, in need of such treatment an effective amount of a compound according to any one of claims 1 to 14.

25. A method for the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa, which comprises administering to a mammal, including humans, in need of such treatment an effective amount of a compound as claimed in any one of claims 1 to 14,
5 wherein the said salt is administered in combination with at least one antimicrobial agent.
26. A pharmaceutical formulation for use in the inhibition of gastric acid secretion wherein the active ingredient is a compound according to any one of claims 1
10 to 14.
27. A pharmaceutical formulation for use in the treatment of gastrointestinal inflammatory diseases wherein the active ingredient is a compound according to any one of claims 1 to 14.
15
28. A pharmaceutical formulation for use in the treatment or prophylaxis of conditions involving infection by *Helicobacter pylori* of human gastric mucosa, wherein the active ingredient is a compound according to any one of claims 1 to 14 in combination with at least one antimicrobial agent.
20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/00991

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 235/08, A61K 31/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CASONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---------------------------------------------------------------------------------------------------------------|-----------------------|
| X | EP 0266326 A1 (AKTIEBOLAGET HÄSSLE), 4 May 1988 (04.05.88) -- | 1-28 |
| X | WO 9604251 A1 (FUJISAWA PHARMACEUTICAL CO., LTD.), 15 February 1996 (15.02.96) -- | 1-28 |
| & | EP 0774462 A1 (FUJISAWA PHARMACEUTICAL CO., LTD.), 21 May 1997 (21.05.97), the claims and page 19 ----- | 1-28 |

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

• Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 October 1997

Date of mailing of the international search report

27 -10- 1997

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Solveig Gustavsson

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 97/00991

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 23-25
because they relate to subject matter not required to be searched by this Authority, namely:

See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

01/10/97

International application No.
PCT/SE 97/00991

| Patent document cited in search report | | | Publication date | Patent family member(s) | | Publication date |
|-------------------------------------------|---------|----|---------------------|----------------------------|------------|---------------------|
| EP | 0266326 | A1 | 04/05/88 | SE | 0266326 T3 | |
| | | | | AU | 7949887 A | 28/04/88 |
| | | | | DE | 3784624 A | 15/04/93 |
| | | | | DK | 555387 D | 00/00/00 |
| | | | | ES | 2053587 T | 01/08/94 |
| | | | | JP | 63122675 A | 26/05/88 |
| | | | | SE | 8604566 D | 00/00/00 |
| | | | | SU | 1597100 A | 30/09/90 |
| | | | | US | 5106862 A | 21/04/92 |
| | | | | ZA | 8707400 A | 27/04/88 |
| | | | | | | |
| WO | 9604251 | A1 | 15/02/96 | AU | 2991595 A | 04/03/96 |
| | | | | EP | 0774462 A | 21/05/97 |
| EP | 0774462 | A1 | 21/05/97 | AU | 2991595 A | 04/03/96 |
| | | | | WO | 9604251 A | 15/02/96 |